

The Evening Sky Map

FREE* EACH MONTH FOR YOU TO EXPLORE, LEARN & ENJOY THE NIGHT SKY

EQUATORIAL EDITION
JULY 2009

Sky Calendar – July 2009

- Earth at Aphelion** (farthest from Sun) at 8h UT. The Sun-Earth distance is 1.01668 a.u. or about 152.1 million km.
- Moon very near Antares** at 9h UT (evening sky). Occultation visible from Japan, the Pacific, and Hawaii.
- Venus near Pleiades** at 14h UT (morning sky).
- Full Moon** at 9:21 UT. The full Moon of July is called the "Thunder Moon" or "Hay Moon".
- Penumbral Eclipse of the Moon** from 8:33 to 10:44 UT, mid-eclipse at 9:39 UT. Too slight to detect visually.
- Moon at apogee** (farthest from Earth) at 22h UT (distance 406,232 km; angular size 29.4').
- Jupiter 0.6° SSE of Neptune** at 17h UT (midnight sky). Mags. -2.7 and +7.8.
- Moon near Jupiter** at 19h UT (midnight sky).
- Mars near Pleiades** at 10h UT (50° from Sun, morning sky). Mag. +1.1.
- Mercury at superior conjunction** with the Sun at 2h UT. The planet passes into the evening sky.
- Venus 3.1° N of Aldebaran** at 6h UT (42° from Sun, morning sky). Mags. -4.1 and +0.9.
- Last Quarter Moon** at 9:53 UT.
- Moon near Pleiades** at 3h UT (morning sky).
- Moon near Mars** at 11h UT (51° from Sun, morning sky).
- Moon near Venus** at 5h UT (41° from Sun, morning sky). Mag. -4.1.
- Moon at perigee** (closest to Earth) at 20h UT (357,463 km; 33.4'). Only 6 hours before New Moon. High tides expected.
- New Moon** at 2:34 UT. Start of lunation 1071.
- Total Solar Eclipse** visible from parts of India, Nepal, China, and central Pacific Ocean. Partial phases visible from most of Asia and Pacific. Greatest totality (6m 39s) occurs at 2:35 UT.
- Moon near Regulus** at 3h UT (evening sky).
- Moon near Saturn** at 10h UT (evening sky). Mag. +1.1.
- First Quarter Moon** at 21:59 UT.
- Moon very near Antares** at 16h UT (evening sky). Occultation visible from S Asia, NE Africa, and SE Europe.


More sky events and links at <http://Skymaps.com/skycalendar/>

All times in Universal Time (UT). (Singapore Standard Time = UT + 8 hours.)

Sky maps Follow us on Twitter
<http://twitter.com/skymaps>



International Year of Astronomy 2009
<http://www.astronomy2009.org>



SKY MAP SHOWS HOW THE NIGHT SKY LOOKS

EARLY JUL 9 PM
LATE JUL 8 PM

SKY MAP DRAWN FOR A LATITUDE OF 0° AND IS SUITABLE FOR LATITUDES UP TO 15° NORTH OR SOUTH OF THIS

Symbols

- Galaxy
- Double Star
- Variable Star
- Diffuse Nebula
- Planetary Nebula
- Open Star Cluster
- Globular Star Cluster

Star Magnitudes -1 0 1 2 3 4

Copyright © 2000–2009 Kym Thalassoudis. All Rights Reserved.

* TERMS OF USE: FREE FOR NON-COMMERCIAL EDUCATIONAL USE. ASTRONOMY EDUCATION GROUPS MAY FREELY DISTRIBUTE PRINTED HANDOUTS. FULL DETAILS AT <http://Skymaps.com/terms.html>

About the Celestial Objects

Listed on this page are several of the brighter, more interesting celestial objects visible in the evening sky this month (refer to the monthly sky map). The objects are grouped into three categories. Those that can be easily seen with the naked eye (that is, without optical aid), those easily seen with binoculars, and those requiring a telescope to be appreciated. **Note, all of the objects (except single stars) will appear more impressive when viewed through a telescope or very large binoculars.** They are grouped in this way to highlight objects that can be seen using the optical equipment that may be available to the star gazer.

Tips for Observing the Night Sky

When observing the night sky, and in particular deep-sky objects such as star clusters, nebulae, and galaxies, it's always best to observe from a dark location. Avoid direct light from street lights and other sources. If possible observe from a dark location away from the light pollution that surrounds many of today's large cities.

You will see more stars after your eyes adapt to the darkness—usually about 10 to 20 minutes after you go outside. Also, if you need to use a torch to view the sky map, cover the light bulb with red cellophane. This will preserve your dark vision.

Finally, even though the Moon is one of the most stunning objects to view through a telescope, its light is so bright that it brightens the sky and makes many of the fainter objects very difficult to see. So try to observe the evening sky on moonless nights around either New Moon or Last Quarter.

Astronomical Glossary

Conjunction – An alignment of two celestial bodies such that they present the least angular separation as viewed from Earth.

Constellation – A defined area of the sky containing a star pattern.

Diffuse Nebula – A cloud of gas illuminated by nearby stars.

Double Star – Two stars that appear close to each other in the sky; either linked by gravity so that they orbit each other (binary star) or lying at different distances from Earth (optical double). Apparent separation of stars is given in seconds of arc (").

Ecliptic – The path of the Sun's center on the celestial sphere as seen from Earth.

Elongation – The angular separation of two celestial bodies. For Mercury and Venus the greatest elongation occurs when they are at their most angular distance from the Sun as viewed from Earth.

Galaxy – A mass of up to several billion stars held together by gravity.

Globular Star Cluster – A ball-shaped group of several thousand old stars.

Light Year (ly) – The distance a beam of light travels at 300,000 km/sec in one year.

Magnitude – The brightness of a celestial object as it appears in the sky.

Open Star Cluster – A group of tens or hundreds of relatively young stars.

Opposition – When a celestial body is opposite the Sun in the sky.

Planetary Nebula – The remnants of a shell of gas blown off by a star.

Universal Time (UT) – A time system used by astronomers. Also known as Greenwich Mean Time. Singapore Standard Time is UT plus 8 hours.

Variable Star – A star that changes brightness over a period of time.

EQUATORIAL EDITION
JULY 2009

CELESTIAL OBJECTS

☆
☆
☆
☆
Sky maps .com

Easily Seen with the Naked Eye

Altair	Aql	• Brightest star in Aquila. Name means "the flying eagle". Dist=16.8 ly.
Arcturus	Boo	• Orange, giant K star. Name means "bear watcher". Dist=36.7 ly.
β Centauri	Cen	• With Alpha Centauri, forms the so-called "Pointers-to-the-Cross". Dist=525 ly.
α Centauri	Cen	• Nearest bright star to Sun at 4.4 ly. Brilliant double star in a telescope. 80 year period.
Coalsack	Cru	♦ Most famous naked-eye dark nebula. Requires dark sky. Dist=600 ly.
α Herculis	Her	⊕ Semi-regular variable. Magnitude varies between 3.1 & 3.9 over 90 days. Mag 5.4 companion.
Vega	Lyr	• The 5th brightest star in the sky. A blue-white star. Dist=25.0 ly.
Antares	Sco	• Red, supergiant star. Name means "rival of Mars". Dist=135.9 ly.
Spica	Vir	• Latin name means "ear of wheat" and shown held in Virgo's left hand. Dist=250 ly.

Easily Seen with Binoculars

η Aquilae	Aql	⊕ Bright Cepheid variable. Mag varies between 3.6 & 4.5 over 7.166 days. Dist=1,200 ly.
6397	Ara	⊕ Thought to be the nearest globular. Dist=7,000 ly.
M3	CVn	⊕ Easy to find in binoculars. Might be glimpsed with the naked eye.
ω Centauri	Cen	⊕ Largest and brightest globular star cluster in sky. 1 million stars. Dist=17,000 ly.
4755	Cru	⊕ Jewel Box. Outstanding star cluster. Many contrasting colours. Dist=7,600 ly.
χ Cygni	Cyg	⊕ Long period pulsating red giant. Magnitude varies between 3.3 & 14.2 over 407 days.
ν Draconis	Dra	• Wide pair of white stars. One of the finest binocular pairs in the sky. Dist=100 ly.
M13	Her	⊕ Best globular in northern skies. Discovered by Halley in 1714. Dist=23,000 ly.
M92	Her	⊕ Fainter and smaller than M13. Use a telescope to resolve its stars.
R Hydrae	Hya	⊕ Long period variable. Mag varies between 3.0 & 11.0 over 390 days. Brilliant red.
ε Lyrae	Lyr	• Famous Double Double. Binoculars show a double star. High power reveals each a double.
R Lyrae	Lyr	⊕ Semi-regular variable. Magnitude varies between 3.9 & 5.0 over 46.0 days.
M12	Oph	⊕ Close to the brighter M10. Dist=18,000 ly.
M10	Oph	⊕ 3 degrees from the fainter M12. Both may be glimpsed in binoculars. Dist=14,000 ly.
IC 4665	Oph	⊕ Large, scattered open cluster. Visible with binoculars.
6633	Oph	⊕ Scattered open cluster. Visible with binoculars.
M8	Sgr	□ Lagoon Nebula. Bright nebula bisected by a dark lane. Dist=5,200 ly.
M25	Sgr	⊕ Bright cluster located about 6 deg N of "teapot's" lid. Dist=1,900 ly.
M22	Sgr	⊕ A spectacular globular star cluster. Telescope will show stars. Dist=10,000 ly.
M4	Sco	⊕ A close globular. May just be visible without optical aid. Dist=7,000 ly.
6231	Sco	⊕ Easy to see in binoculars. Dist=5,900 ly.
M6	Sco	⊕ Butterfly Cluster. 30+ stars in 7x binoculars. Dist=1,960 ly.
M7	Sco	⊕ Superb open cluster. Visible to the naked eye. Age=260 million years. Dist=780 ly.
M5	Ser	⊕ Fine globular star cluster. Telescope will reveal individual stars. Dist=25,000 ly.
6025	TrA	⊕ A small open star cluster in Milky Way. Dist=2,700 ly.
Cr 399	Vul	⊕ Coathanger asterism or "Brocchi's Cluster". Not a true star cluster. Dist=218 to 1,140 ly.

Telescopic Objects

ε Boötis	Boo	• Red giant star (mag 2.5) with a blue-green mag 4.9 companion. Sep=2.8". Difficult to split.
5128	Cen	◊ Bisected by a wide obscuring lane. Strong radio source. Dist=11 million ly.
M64	Com	◊ Black-Eye Galaxy. Discovered by J.E. Bode in 1775 -- "a small, nebulous star".
Albireo	Cyg	• Beautiful double star. Contrasting colours of orange and blue-green. Sep=34.4".
M83	Hya	◊ Classic face-on spiral. Discovered in 1752 by Lacaille. In attractive star field.
5822	Lup	◊ Large, attractive cluster. Dist=1,800 ly. Open cluster NGC 5823 to the south.
β Lyrae	Lyr	⊕ Eclipsing binary. Mag varies between 3.3 & 4.3 over 12.940 days. Fainter mag 7.2 blue star.
M57	Lyr	♦ Ring Nebula. Magnificent object. Smoke-ring shape. Dist=4,100 ly.
M23	Sgr	◊ Elongated star cluster. Telescope required to show stars. Dist=2,100 ly.
M20	Sgr	□ Trifid Nebula. A telescope shows 3 dust lanes trisecting nebula. Dist=5,200 ly.
M21	Sgr	⊕ A fine and impressive cluster. Dist=4,200 ly.
M17	Sgr	□ Omega Nebula. Contains the star cluster NGC 6618. Dist=4,900 ly.
6124	Sco	◊ Contains 5 bright tightly packed stars near centre. 7 star chain. Dist=1,600 ly.
M11	Sct	◊ Wild Duck Cluster. Resembles a globular through binoculars. V-shaped. Dist=5,600 ly.
M16	Ser	◊ Eagle Nebula. Requires a telescope of large aperture. Dist=8,150 ly.
M104	Vir	◊ Sombrero Galaxy. Almost edge-on spiral galaxy. Protruding central core.
γ Virginis	Vir	• Superb pair of mag 3.5 yellow-white stars. Orbit=169 years. At their closest in 2005.
M27	Vul	♦ Dumbbell Nebula. Large, twin-lobed shape. Most spectacular planetary. Dist=975 ly.